

Synonym

MSLN, Mesothelin, MPF

Source

PE-Labeled Human Mesothelin (296-580), His Tag (MSN-HP223) is expressed from human HEK293 cells. It contains AA Glu 296 - Gly 580 (Accession # [AAH09272](#)). It is the PE labeled form of Human Mesothelin (296-580), His Tag. Predicted N-terminus: Glu 296

Molecular Characterization

Mesothelin(Glu 296 - Gly 580) AAH09272	Poly-his
---	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 33.0 kDa.

Conjugate

PE

Excitation Wavelength: 488 nm / 561 nm

Emission Wavelength: 575 nm

Formulation

Lyophilized from 0.22 μ m filtered solution in 10 mM Tris, 150 mM NaCl, 0.1% BSA, pH8.2. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

Bioactivity-FACS

5e5 of anti-MSLN CAR-293 cells were stained with 100 μ L of 1:25 dilution (4 μ L stock solution in 100 μ L FACS buffer) of PE-Labeled Human Mesothelin (296-580), His Tag (Cat. No. MSN-HP223) and negative control protein respectively. PE signal was used to evaluate the binding activity (QC tested).

Background

Mesothelin (MSLN) is also known as CAK1 antigen, Pre-pro-megakaryocyte-potentiating factor, which belongs to the mesothelin family. Mesothelin / MSLN can be proteolytically cleaved into the following two chains by a furin-like convertase: Megakaryocyte-potentiating factor (MPF) and the cleaved form of mesothelin. Both

MPF and the cleaved form of mesothelin are N-glycosylated. Mesothelin / MSLN can interact with MUC16. The membrane-anchored forms of MSLN may play a role in cellular adhesion. MPF potentiates megakaryocyte colony formation in vitro.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.