

Synonym

OX40L, TNFSF4, CD252, Glycoprotein Gp34, TXGP1, CD134 ligand, CD134L

Source

Human OX40 Ligand, Fc Tag (OXL-H526x) is expressed from human 293 cells (HEK293). It contains AA Gln 51 - Leu 183 (Accession # [NP_003317](#)).

Molecular Characterization

Fc(Thr 106 - Lys 330) P01857	OX40 Ligand(Gln 51 - Leu 183) NP_003317
---------------------------------	--

This protein carries a human IgG1 Fc tag at the N-terminus.

The protein has a calculated MW of 42.4 kDa. The protein migrates as 53-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Human OX40 Ligand, Fc Tag on SDS-PAGE under reducing (R) condition.


The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA




Immobilized Human OX40, His Tag (Cat. No. [OX0-H5224](#)) at 0.5 µg/mL (100 µL/well) can bind Human OX40 Ligand, Fc Tag (Cat. No. [OXL-H526x](#)) with a linear range of 2-63 ng/mL (QC tested).

Bioactivity-BLI



Loaded Human OX40 Ligand, Fc Tag (Cat. No. [OXL-H526x](#)) on Protein A Biosensor, can bind Human OX40, His Tag (Cat. No. [OX0-H5224](#)) with an affinity constant of 0.96 µM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Bioactivity-FACS



FACS assay shows that recombinant Human OX40 Ligand, Fc Tag (Cat. No. OXL-H526x) can bind to 293T cell overexpressing human OX40. The concentration of OX40 Ligand used is 0.01 µg/ml (Routinely tested).

Background

Tumor necrosis factor ligand superfamily member 4 (TNFSF4) is also known as glycoprotein Gp34, OX40 ligand (OX40L), TAX transcriptionally-activated glycoprotein 1 and CD252, which belongs to the tumor necrosis factor family. TNFSF4 is the ligand for CD134 and is expressed on such cells as DC2s (a subtype of dendritic cells) enabling amplification of Th2 cell differentiation. The interaction of TNFSF4-TNFSF4 is involved in the pathogenesis of multiple autoimmune and inflammatory diseases such as systemic lupus erythematosus (SLE), carotid artery disease and cancer. Furthermore, similar to other TNF superfamily members, membrane-bound OX40 Ligand (TNFSF4) exists as a homotrimer. Human TNFSF4 shares 46% amino acid sequence identity with its mouse counterpart.

References

- (1) [Compaan D.M., et al., 2006, Structure 14:1321-1330.](#)
- (2) [Maizels R.M., et al., 2003, Nat. Rev. Immunol. 3 \(9\): 733-44.](#)

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.