

Synonym

TNFRSF9,4-1BB,CD137,CDw137,ILA

Source

Human 4-1BB Protein, Mouse IgG2a Fc Tag(41B-H5256) is expressed from human 293 cells (HEK293). It contains AA Leu 24 - Gln 186 (Accession # [Q07011-1](#) ).

Predicted N-terminus: Leu 24

Molecular Characterization

4-1BB(Leu 24 - Gln 186) Q07011-1	mFc(Glu 98 - Lys 330) P01863
-------------------------------------	---------------------------------

This protein carries a mouse IgG2a Fc tag at the C-terminus

The protein has a calculated MW of 44.2 kDa. The protein migrates as 55-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:



- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Human 4-1BB Protein, Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA




Immobilized Human 4-1BB Ligand, Fc Tag (Cat. No. 41L-H5257) at 1 µg/mL (100 µL/well) can bind Human 4-1BB Protein, Mouse IgG2a Fc Tag (Cat. No. 41B-H5256) with a linear range of 0.6-5 ng/mL (QC tested).

Immobilized Human 4-1BB Ligand, His Tag (Cat. No. 41L-H5249) at 2 µg/mL (100 µL/well) can bind Human 4-1BB Protein, Mouse IgG2a Fc Tag (Cat. No. 41B-H5256) with a linear range of 0.1-4 ng/mL (Routinely tested).



Immobilized Human 4-1BB Protein, Mouse IgG2a Fc Tag (Cat. No. 41B-H5256) at 1 µg/mL (100 µL/well) can bind Anti-Human 4-1BB MAb, Human IgG4 with a linear range of 0.2-6 ng/mL (Routinely tested).

Bioactivity-FACS



Flow Cytometry assay shows that Human 4-1BB Protein, Mouse IgG2a Fc Tag (Cat. No. 41B-H5256) can bind to 293T cells overexpressing Human 4-1BBL. The concentration of 4-1BB used is 0.03 µg/mL (Routinely tested).

## Background

4-1BB is also known as CD137, tumor necrosis factor receptor superfamily member 9 (TNFRSF9), induced by lymphocyte activation (ILA), is a co-stimulatory molecule of the tumor necrosis factor (TNF) receptor superfamily. CD137 can be expressed by activated T cells, but to a larger extent on CD8 than on CD4 T cells. In addition, CD137 expression is found on dendritic cells, follicular dendritic cells, natural killer cells, granulocytes and cells of blood vessel walls at sites of inflammation. The best characterized activity of CD137 is its costimulatory activity for activated T cells. Crosslinking of CD137 enhances T cell proliferation, IL-2 secretion survival and cytolytic activity. Further, it can enhance immune activity to eliminate tumors in mice. CD137 can enhance activation-induced T cell apoptosis when triggered by engagement of the TCR/CD3 complex. In addition, 4-1BB/4-1BBL co-stimulatory pathway has been shown to augment secondary CTL responses to several viruses, and meanwhile augment anti-tumor immunity. 4-1BB thus is a promising candidate for immunotherapy of human cancer. CD137 has been shown to interact with TRAF2.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.