

Synonym

Anterior gradient protein 2 homolog,AG-2,hAG-2,HPC8,Secreted cement gland protein XAG-2 homolog,AGR2

Source

Mouse AGR2, His Tag (AG2-M52H6) is expressed from human 293 cells (HEK293). It contains AA Lys 21 - Leu 175 (Accession # [O88312-1](#)).
Predicted N-terminus: Lys 21

Molecular Characterization

AGR2(Lys 21 - Leu 175)
O88312-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.
The protein has a calculated MW of 19.8 kDa. The protein migrates as 23 kDa and 24 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.
>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, 1 mM TCEP, pH7.4.
Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:
• -20°C to -70°C for 12 months in lyophilized state;
• -70 °C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse AGR2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS



The purity of Mouse AGR2, His Tag (Cat. No. AG2-M52H6) was more than 90% and the molecular weight of this protein is around 55-68kDa verified by SEC-MALS.
[Report](#)

Background

Anterior gradient 2 (AGR2) is a normal endoplasmic reticulum protein that has two important abnormal functions, amphibian limb regeneration and human cancer metastasis promotion. These normal intracellular and abnormal extracellular roles can be attributed to the multidomain structure of AGR2. The NMR structure shows that AGR2 consists of an unstructured N-terminal region followed by a thioredoxin fold.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.