Biotinylated Mouse BAFF / TNFSF13B / CD257 Protein, Avitag™,Fc Tag, active trimer (MALS verified)

Catalog # BAF-M82F7



Synonym

TNFSF13B,BAFF,BLYS,CD257,DTL,TALL1,THANK,TNFSF20,ZTNF4,TALL-1

Source

Biotinylated Mouse BAFF Protein, Avitag,Fc Tag(BAF-M82F7) is expressed from human 293 cells (HEK293). It contains AA Arg164 - Leu309 (Accession # Q9WU72-1).

Predicted N-terminus: Gly

Molecular Characterization



This protein carries an Avi tag (AvitagTM) at the N-terminus, followed by a human IgG1 Fc tag

The protein has a calculated MW of 44.8 kDa. The protein migrates as 50-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using AvitagTM technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from $0.22~\mu m$ filtered solution in 50~mM Tris, 100~mM Glycine, 150~mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Mouse BAFF Protein, Avitag,Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA

SEC-MALS




The purity of Biotinylated Mouse BAFF Protein, Avitag,Fc Tag (Cat. No. BAF-M82F7) is more than 95% and the molecular weight of this protein is around 265-290 kDa verified by SEC-MALS.

Report

Biotinylated Mouse BAFF / TNFSF13B / CD257 Protein, Avitag™,Fc Tag, active trimer (MALS verified)

Catalog # BAF-M82F7





Immobilized Mouse BAFFR, His Tag (Cat. No. BAR-M52H8) at 2 μ g/mL (100 μ L/well) can bind Biotinylated Mouse BAFF Protein, Avitag,Fc Tag (Cat. No. BAF-M82F7) with a linear range of 0.2-8 μ g/mL (QC tested).

Background

B-cell activating factor (BAFF) is also known as tumor necrosis factor ligand superfamily member 13B, TNFSF13B, BAFF, B Lymphocyte Stimulator (BLyS), cluster of differentiation 257 (CD257), DTL, TNF- and APOL-related leukocyte expressed ligand (TALL-1), THANK, TNFSF20, ZTNF4, and is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptors TNFRSF13B/TACI, TNFRSF17/BCMA, and TNFRSF13C/BAFFR. This cytokine is expressed in B cell lineage cells, and acts as a potent B cell activator. It has been also shown to play an important role in the proliferation and differentiation of B cells. It is expressed as transmembrane protein on various cell types including monocytes, dendritic cells and bone marrow stromal cells. BAFF is the natural ligand of three unusual tumor necrosis factor receptors named BAFF-R, TACI, and BCMA, all of which have differing binding affinities for it. These receptors are expressed mainly on mature B lymphocytes (TACI is also found on a subset of T-cells and BCMA on plasma cells). TACI binds worst since its affinity is higher for a protein similar to BAFF, called a proliferation-inducing ligand (APRIL). BCMA displays an intermediate binding phenotype and will work with either BAFF or APRIL to varying degrees. Signaling through BAFF-R and BCMA stimulates B lymphocytes to undergo proliferation and to counter apoptosis. All these ligands act as heterotrimers (i.e. three of the same molecule) interacting with heterotrimeric receptors, although BAFF has been known to be active as either a heteroor homotrimer. BAFF acts as a potent B cell activator and has been shown to play an important role in the proliferation and differentiation of B cells.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.