

Synonym

CCR5,CMKBR5,CD195,CC-CKR-5,CHEMR13

Source

Human CCR5, Flag,His Tag(CC5-H52D1) is expressed from human 293 cells (HEK293). It contains AA Asp 2 - Arg 223 & Glu 227 - Arg 319 (Accession # P51681-1 (C58Y, G163N, A233D, K303E)).

Predicted N-terminus: Asp

Molecular Characterization



This protein carries flag tag at the N-terminus and polyhistidine tag at the C-terminus.

The protein has a calculated MW of 46.0 kDa. The protein migrates as 40-43 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>85% as determined by SDS-PAGE.

Formulation

This product is not suitable for cell based experiments due to cytotoxicity of DDM.

DDM and CHS are INDISPENSABLE to keep membrane protein soluble and active, under no circumastance should you remove DDM and CHS.

DDM/CHS buffer (DC-11) is sold separately and not included in protein, and please contact us if you need the buffer.

If glycerol is not compatible to your application, remove glycerol just before immediate experiment, and NEVER store glycerol-free protein solution.

Supplied as 0.2 µm filtered solution in 50 mM HEPES, 150 mM NaCl, DDM, CHS, pH7.5 with glycerol as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping cost.


Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.

SDS-PAGE




Human CCR5, Flag, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 85% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

Bioactivity-ELISA

Discounts, Gifts, and more!


^{**}The DDM/CHS buffer (Cat. No. DC-11) is sold separately and not included in protein, you can follow this link for product information.






Immobilized Human CCR5, Flag,His Tag (Cat. No. CC5-H52D1) at $10 \mu g/mL$ (100 $\mu L/well$) can bind Anti-Human CCR5 Antibody with a linear range of 0.02-1.25 $\mu g/mL$ (QC tested).

Bioactivity-SPR



Human CCR5, Flag, His Tag (Cat. No. CC5-H52D1) captured on CM5 chip via anti-His antibody can bind Anti-CCR5 antibody (Human IgG1) with an affinity constant of 0.0737 nM as determined in a SPR assay (in presence of DDM and CHS) (Biacore T200) (Routinely tested).



Purified NA/LE Mouse Anti-Human CD195 (Clone: 2D7) (Mouse IgG2a) captured on CM5 chip via anti-mouse antibodies surface can bind Human CCR5, Flag,His Tag (Cat. No. CC5-H52D1) with an affinity constant of 7.51 nM as determined in a SPR assay (in presence of DDM and CHS) (Biacore T200) (Routinely tested).

Background

Receptor for a number of inflammatory CC-chemokines including CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation. Publications (Microbial infection) Acts as a coreceptor (CD4 being the primary receptor) of human immunodeficiency virus-1/HIV-1.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.

