

## **Synonym**

MMR,CD206,hMR,MRC1,CLEC13D,CLEC13DL,MRC1L1

#### Source

Human CD206, His Tag(CD6-H52H9) is expressed from human 293 cells (HEK293). It contains AA Leu 19 - Ala 1389 (Accession # P22897-1). Predicted N-terminus: Leu 19

### **Molecular Characterization**

CD206(Leu 19 - Ala 1389) P22897-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 158.6 kDa. The protein migrates as 150-190 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### Endotoxin

Less than 1.0 EU per µg by the LAL method.

## **Purity**

>90% as determined by SDS-PAGE.

#### **Formulation**

Lyophilized from 0.22  $\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## **SDS-PAGE**



Human CD206, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

# **Background**

Mrc1 is a conserved checkpoint mediator protein that transduces the replication stress signal to the downstream effector kinase. Mrc1 and its vertebrate homologue Claspin serve as a mediator for replication stress checkpoint signaling, receiving the signal from Mec1/Rad3/ATR sensor kinase and transmitting it to the effector Rad53/Cds1/Chk1 kinase. The loss of mrc1 checkpoint activity results in the aberrant activation of late/dormant origins in the presence of hydroxyurea. Tumor-



# **Human CD206 / MMR Protein, His Tag**





associated macrophages (TAMs) expressing the multi-ligand endocytic receptor mannose receptor (CD206/MRC1) contribute to tumor immunosuppression, angiogenesis, metastasis, and relapse.

**Clinical and Translational Updates** 

