Catalog # CD7-H82F3



Synonym

CD7,GP40,TP41,LEU-9,Tp40

Source

Biotinylated Human CD7, Fc, Avitag(CD7-H82F3) is expressed from human 293 cells (HEK293). It contains AA Ala 26 - Pro 180 (Accession # <u>P09564-1</u>). Predicted N-terminus: Ala 26

Molecular Characterization



This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (AvitagTM).

The protein has a calculated MW of 44.6 kDa. The protein migrates as 55-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag[™] technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per μ g by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Human CD7, Fc, Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS



The purity of Biotinylated Human CD7, Fc, Avitag (Cat. No. CD7-H82F3) is more than 90% and the molecular weight of this protein is around 95-115 kDa verified by SEC-MALS.



Bioactivity-ELISA




>>> www.acrobiosystems.com



Catalog # CD7-H82F3





Immobilized Biotinylated Human CD7, Fc, Avitag (Cat. No. CD7-H82F3) at 1 μ g/mL (100 μ L/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 μ g/well) plate can bind Anti-CD7 antibody, Mouse IgG1 with a linear range of 0.2-2 ng/mL (QC tested).

Background

T-cell antigen CD7 (CD7) is also known as GP40, LEU-9, TP41 and Tp40. CD7 is a protein that in humans is encoded by the CD7 gene, this gene encodes a transmembrane protein which is a member of the immunoglobulin superfamily. CD7 has been shown to interact with PIK3R1. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.



>>> www.acrobiosystems.com

