


Synonym

ERBB2,CD340,HER-2/neu,HER2,MLN19,NEU,NGL,TKR1

Source

Mouse Her2, His Tag (ER2-M5220) is expressed from human 293 cells (HEK293). It contains AA Thr 23 - Thr 653 (Accession # AAH53078.1).  
Predicted N-terminus: Thr 23

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 70.6 kDa. The protein migrates as 80-95 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

- This product is stable after storage at:
- -20°C to -70°C for 12 months in lyophilized state;
  - -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse Her2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Background

Human Epidermal growth factor Receptor 2 (HER2) is also called ERBB2, HER-2,HER-2 /neu, NEU, NGL,TKR1 and c-erb B2,and is a protein giving higher aggressiveness in breast cancers. It is a member of the ErbB protein family, more commonly known as the epidermal growth factor receptor family. HER2 is a cell membrane surface-bound receptor tyrosine kinase and is normally involved in the signal transduction pathways leading to cell growth and differentiation. HER2 is thought to be an orphan receptor, with none of the EGF family of ligands able to activate it. Approximately 30% of breast cancers have an amplification of the HER2 gene or overexpression of its protein product. Overexpression of this receptor in breast cancer is associated with increased disease recurrence and worse prognosis. HER2 appears to play roles in development, cancer, communication at the neuromuscular junction and regulation of cell growth and differentiation .

References

(1) [Kanai, Y. et. al., 1995, Biochem. Biophys. Res. Commun.208:1067.](#)  
(2) [Codony-Seeervat, J. et al., 1999, Cancer Res. 59: 1196-1201.](#)  
(3) [Tzahar, E. et al., 1998, Biochim. Biophys. Acta. 1377: 25-37.](#)  
(4) [Kermit, L. et al, 1999, J. Biol. Chem. 274: 5263-5266.](#)  
(5) [Hellyer, N.J. et. al., 2001, J. Biol. Chem. 276:42153.](#)  
(6) [Daly, R.J., 1999, Growth Factors 16:255.](#)

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.