

Synonym

FAP,FAPalpha,SIMP,Seprase,APCE

Source

Human FAP, Fc Tag(FAP-H5263) is expressed from human 293 cells (HEK293). It contains AA Leu 26 - Asp 760 (Accession # [Q12884-1](#)).
Predicted N-terminus: Pro

Molecular Characterization

Fc(Pro 100 - Lys 330)	FAP(Leu 26 - Asp 760)
P01857	Q12884-1

This protein carries a human IgG1 Fc tag at the N-terminus.
The protein has a calculated MW of 111.5 kDa. The protein migrates as 120-130 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.
>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.
This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human FAP, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS




The purity of Human FAP, Fc Tag (Cat. No. FAP-H5263) is more than 90% and the molecular weight of this protein is around 216-264 kDa verified by SEC-MALS.
[Report](#)



Human FAP Protein, Fc Tag, active dimer (MALS verified)


Catalog # FAP-H5263



Immobilized Human FAP, Fc Tag (Cat. No. FAP-H5263) at 1 µg/mL (100 µL/well) can bind Anti-FAP Antibody, Mouse IgG2a (FAP5) with a linear range of 0.2-3 ng/mL (QC tested).

Bioactivity

Measured by its ability to convert the substrate benzyloxycarbonyl-Gly-Pro-7-amido-4-methylcoumarin (Z-GP-AMC) to Z-Gly-Pro and 7-amino-4-methylcoumarin (AMC). The specific activity is >7000 pmol/min/µg (QC tested).



Immobilized Anti-FAP Antibody, Mouse IgG2a (FAP5) at 1 µg/mL (100 µL/well) can bind Human FAP, Fc Tag (Cat. No. FAP-H5263) with a linear range of 0.5-4 ng/mL (Routinely tested).

Background

FAP (also known as seprase) is a Type II transmembrane serine protease. Both plasma membrane and soluble forms exhibit post-proline cleaving endopeptidase activity, with a marked preference for Ala/Ser-Gly-Pro-Ser/Asn/Ala consensus sequences. Degrade also gelatin, heat-denatured type I collagen. Also has dipeptidyl peptidase activity, with a preference for Ala-Pro, Ile-Pro, Gly-Pro, Arg-Pro and Pro-Pro. The plasma membrane form, in association with either DPP4, PLAUR or integrins, is involved in the pericellular proteolysis of the extracellular matrix (ECM), and hence promotes cell adhesion, migration and invasion through the ECM. Promotes glioma cell invasion through the brain parenchyma by degrading the proteoglycan brevican. Acts as a tumor suppressor in melanocytic cells through regulation of cell proliferation and survival in a serine protease activity-independent manner.

Clinical and Translational Updates

