# Biotinylated Canine FcRn / FCGRT&B2M Heterodimer Protein, His,Avitag™&Tag Free (MALS & SPR verified)

Catalaa # FCN/ C00\//0



#### Synonym


FcRn,FCGRT & B2M

#### Source

Biotinylated Canine FCGRT&B2M Heterodimer Protein, His,Avitag&Tag Free(FCM-C82W9) is expressed from human 293 cells (HEK293). It contains AA Ala 23 - Pro 298 & Val 28 - Asn 125 (Accession # XP\_038512242.1 & NP\_001271408.1).

Predicted N-terminus: Ala 23 | Val 28

#### **Molecular Characterization**



This protein carries a polyhistidine tag at the N-terminus. The protein has a calculated MW of  $28.3 \text{ kDa} \mid 11.5 \text{ kDa}$ . The protein migrates as  $\leq 14.4 \text{ kDa}$  and 38-40 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

## Labeling

Biotinylation of this product is performed using Avitag<sup>TM</sup> technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

### **Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

## **Endotoxin**

Less than  $1.0 \ EU$  per  $\mu g$  by the LAL method.

### **Purity**

>95% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

#### **Formulation**

Lyophilized from  $0.22~\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

### **Storage**


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


## **SDS-PAGE**



Biotinylated Canine FCGRT&B2M Heterodimer Protein, His,Avitag&Tag Free on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

# **Bioactivity-ELISA**

## **SEC-MALS**




The purity of Biotinylated Canine FCGRT&B2M Heterodimer Protein, His,Avitag&Tag Free (Cat. No. FCM-C82W9) is more than 95% and the molecular weight of this protein is around 43-53 kDa verified by SEC-MALS. Report

# Biotinylated Canine FcRn / FCGRT&B2M Heterodimer Protein, His,Avitag™&Tag Free (MALS & SPR verified)



Catalaa # FONA COONA


Biotinylated Canine FcRn / FCGRT&B2M Heterodimer, His,Avitag&Tag Free ELISA 0.1 µg of Monoclonal Anti-Human CTLA-4 Antibody, Human IgG1 per well




Biotinylated Canine FcRn / FCGRT&B2M Heterodimer, His, Avitag&Tag Free Conc. (μg/mL)

Immobilized Monoclonal Anti-Human CTLA-4 Antibody, Human IgG1 at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Biotinylated Canine FcRn / FCGRT&B2M Heterodimer, His,Avitag&Tag Free (Cat. No. FCM-C82W9) with a linear range of 0.039-2.5  $\mu$ g/mL (Routinely tested).


## **Bioactivity-SPR**



Biotinylated Canine FCGRT&B2M Heterodimer, His,Avitag&Tag Free (Cat. No. FCM-C82W9) immobilized on SA Chip can bind Ipilimumab with an affinity constant of 42.5 nM as determined in a SPR assay (Biacore 8K) (QC tested).



Biotinylated Canine FCGRT&B2M Heterodimer, His,Avitag&Tag Free (Cat. No. FCM-C82W9) immobilized on SA Chip can bind Trastuzumab with an affinity constant of 44.6 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).



Biotinylated Canine FCGRT&B2M Heterodimer, His, Avitag&Tag Free (Cat. No. FCM-C82W9) captured on CM5 chip via anti-His antibody can bind Ipilimumab with an affinity constant of 0.267 μM as determined in a SPR assay (Biacore 8K) (Routinely tested).

# Biotinylated Canine FcRn / FCGRT&B2M Heterodimer Protein, His,Avitag™&Tag Free (MALS & SPR verified)

ACCO

Catalaa # FONA COONAO

## **Background**

FCGRT & B2M heterodimer protein (FcRn complex) consist of two subunits: p51 (equivalent to FCGRT), and p14 (equivalent to beta-2-microglobulin), and forms an MHC class I-like heterodimer. Fc fragment of IgG, receptor, transporter, alpha (FCGRT) binds to the Fc region of monomeric immunoglobulins gamma and mediates the uptake of IgG from milk. FCGRT possible role in transfer of immunoglobulin G from mother to fetus. Beta-2-microglobulin (B2M) is a component of the class I major histocompatibility complex (MHC) and involved in the presentation of peptide antigens to the immune system.

## **Clinical and Translational Updates**

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.