

**Synonym**

gastrin-17,gastrin 17,gastrin

**Source**

Biotinylated Human Gastrin-17, Fc,Avitag (GA7-H82F4) is expressed from human 293 cells (HEK293).

**Molecular Characterization**

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™)

The protein has a calculated MW of 30.3 kDa. The protein migrates as 34-40 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Labeling**

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

**Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

**Endotoxin**

Less than 0.1 EU per µg by the LAL method.

**Sterility**

The sterility testing was performed by membrane filtration method.

**Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

**Storage**


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


**SDS-PAGE**



Biotinylated Human Gastrin-17, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.


**Bioactivity-ELISA**

**SEC-MALS**



The purity of Biotinylated Human Gastrin-17, Fc,Avitag (Cat. No. GA7-H82F4) is more than 90% and the molecular weight of this protein is around 60-70 kDa verified by SEC-MALS.

[Report](#)



Immobilized Anti-Gastrin-17 antibody, Human IgG1 at 2 µg/mL (100 µL/well) can bind Biotinylated Human Gastrin-17, Fc,Avitag (Cat. No. GA7-H82F4) with a linear range of 0.1-2.5 ng/mL (QC tested).

## Background

Gastrin is a peptide hormone that stimulates secretion of gastric acid (HCl) by the parietal cells of the stomach and aids in gastric motility. It is released by G cells in the pyloric antrum of the stomach, duodenum, and the pancreas. Gastrin binds to cholecystinin B receptors to stimulate the release of histamines in enterochromaffin-like cells, and it induces the insertion of K<sup>+</sup>/H<sup>+</sup> ATPase pumps into the apical membrane of parietal cells (which in turn increases H<sup>+</sup> release into the stomach cavity). Its release is stimulated by peptides in the lumen of the stomach.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.