

Synonym

HA, Hemagglutinin

Source

Influenza A (Guangdong/18SF020(H5N6)) Hemagglutinin (HA) Protein, His Tag (HA6-V52H3) is expressed from human 293 cells (HEK293). It contains AA Asp 11 - Ser 526 (Accession # EPI1366513, GISAID).

Predicted N-terminus: Asp 11

Molecular Characterization

HA(Asp 11 - Ser 526) EPI1366513

Poly-his

This protein carries a polyhistidine tag at the C-terminus

The protein has a calculated MW of 63.8 kDa. The protein migrates as 80-95 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Influenza A (Guangdong/18SF020(H5N6)) Hemagglutinin (HA) Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With Star Ribbon Pre-stained Protein Marker).


Bioactivity-ELISA

Influenza A (Guangdong/18SF020(H5N6)) Hemagglutinin (HA) Protein, His Tag





Influenza A (Guangdong/18SF020(H5N6)) Hemagglutinin (HA) Protein, His Tag ELISA 0.1 μ g of Influenza A (Guangdong/18SF020(H5N6)) Hemagglutinin (HA) Protein, His Tag per well



Immobilized Influenza A (Guangdong/18SF020(H5N6)) Hemagglutinin (HA) Protein, His Tag (Cat. No. HA6-V52H3) at 1 μ g/mL (100 μ L/well) can bind Anti-HACL:FC41 Human IgG1 with a linear range of 0.1-4 ng/mL (QC tested).

Background

Neuraminidase (NA) and hemagglutinin (HA) are major membrane glycoproteins found on the surface of influenza virus. Hemagglutinin binds to the sialic acid-containing receptors on the surface of host cells during initial infection and at the end of an infectious cycle. Hemagglutinin also plays a major role in the determination of host range restriction and virulence. As a class I viral fusion protein, hemagglutinin is responsible for penetration of the virus into the cell cytoplasm by mediating the fusion of the membrane of the endocytosed virus particle with the endosomal membrane.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.