



# Background

The Claudin-18.2 (CLDN 18.2) protein, an isoform of Claudin 18, a member of the tight junction protein family, is a highly selective biomarker with limited expression in normal tissues and often abnormal expression during the occurrence and development of various primary malignant tumors, such as gastric cancer/gastroesophageal junction (GC/GEJ) cancer, breast cancer, colon cancer, liver cancer, head and neck cancer, bronchial cancer and non-small-cell lung cancer. Claudin-18.2 participates in the proliferation, differentiation and migration of tumor cells. Recent studies have identified Claudin-18.2 expression as a potential specific marker for the diagnosis and treatment of these tumors.

#### **Host Species**

Mouse

Clone

3B10

**Application** 

IHC

**Property** 

1: 1000

State

Liquid

#### **Positive Control**

**Human Stomach Tissues** 

**Clonality** 

Monoclonal

Synonym

Claudin18.2, CLDN 18.2

**Research Field** 

**Cancer Drug Targets** 

Source

Mouse

**Isotype** 

IgG

Storage


Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.

### **Typical Data**

# **Control Sample**



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278) Human Stomach Tissue, 4X




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278)


Human Stomach Tissue, 20X

Immunohistochemical analysis of paraffin embedded human stomach tissue labelled with HCS-S278 at 1/1000 dilution. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.






Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278) Human Lung Tissue, 4X




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278) Human Lung Tissue, 20X

Immunohistochemical analysis of paraffin-embedded human lung tissue labeling Claudin-18.2 with HCS-S278 at 1/1000 dilution, followed by Goat Anti-Mouse IgG H&L (HRP) ready to use. The staining in the lung tissue sample was negative. Counter stained with Hematoxylin. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

#### **Cancer Sample**




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278)




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278)

Human Colorectal Cancer, 20X



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278) Human Pancreatic Cancer, 20X



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) -Recombinant Monoclonal Anti-Claudin-18.2 Antibody, Mouse (3B10) (HCS-S278)

Human Ovarian Cancer, 20X

Immunohistochemical analysis of paraffin-embedded human cancer tissue labeling Claudin-18.2 with HCS-S278 at 1/1000 dilution, followed by Goat Anti-Mouse IgG H&L (HRP) ready to use. Membranous staining on tumor cells is observed. Counter stained with Hematoxylin. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

### Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.

