
Catalog # HGS-S263



Background	Positive Control
Involved in T-cell adhesion processes and in spontaneous rosette formation with	Human Tonsil Tissues
erythrocytes. Plays a role in a late step of leukocyte extravasation helping	Clonality
leukocytes to overcome the endothelial basement membrane. Acts at the same site as, but independently of, PECAM1. Involved in T-cell adhesion processes	Monoclonal
(By similarity).	Synonym
Host Species	MIC2
Rabbit	Research Field
Clone	Cancer Drug Targets
4F5	Source
Application	Rabbit
Аррисации	Isotype
IHC	IgG
Property	Storage
1: 1000	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot.
State	Store at -20°C long term. Avoid freeze / thaw cycle.
Liquid	

## **Typical Data**

**Control Sample** 



Human Tonsil Tissue, 20X

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-CD99 Antibody, Rabbit (4F5) (HGS-S263) Human Tonsil Tissue, 20X

Immunohistochemical analysis of paraffin embedded Human tonsil tissue labelled with HGS-S263 at 1/1000 dilution. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

**Cancer Sample** 




>>> www.acrobiosystems.com



## Recombinant Monoclonal Anti-CD99 Antibody, Rabbit (4F5)

Catalog # HGS-S263





Immunohistochemical analysis of paraffin embedded human cancer tissue labelled with HGS-S263 at 1/1000 dilution. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

**Clinical and Translational Updates** 

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.



>>> www.acrobiosystems.com

