

Source

HRP conjugated Anti-Human-IgG-Fc Antibody (6F11C8), mAb antibody is produced from a hybridoma resulting from fusion of SP2/0 myeloma and B-lymphocytes obtained from a mouse immunized with IgG.

Isotype

Mouse IgG1 | Mouse Kappa

Specificity

This product is a specific antibody specifically reacts with IgG.

Application

ELISA, WB

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH 7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 6 months after reconstitution;
- $2-8^{\circ}\text{C}$ for 2-3 weeks under sterile conditions after reconstitution.

Bioactivity-Elisa

HRP conjugated Anti-Human-IgG-Fc Antibody (6F11C8),mAb ELISA 0.2ug of Human TNF-alpha Protein, His Tag per well



Immobilized Human TNF-alpha Protein, His Tag (Cat. No. TNA-H5228) at $2\ \mu\text{g}/\text{mL}$ ($100\ \mu\text{L}/\text{well}$) can bind Human Monoclonal Anti-TNF-alpha antibody, Human IgG1 (16H5) (Cat. No. TNA-AM494) when detected by HRP conjugated Anti-Human-IgG-Fc Antibody (6F11C8),mAb (Cat. No. IGG-LY69) dilute at 1:10000 ($0.0842\ \mu\text{g}/\text{ml}$) (QC tested).

Background

Immunoglobulins can be divided into five main classes/isotypes which are IgA, IgD, IgE, IgG, and IgM. IgG class identity is determined by class-specific sequences in the Fc region of the heavy chain. IgG antibody class are the most abundant immunoglobulins isotype in blood, lymph fluid, cerebrospinal fluid and peritoneal

fluid. IgGs include four subclasses (IgG1, IgG2, IgG3, and IgG4). The IgG subclasses differ in their physical and chemical properties. Their distribution pattern is found to be age dependent and every subclass has a specific biological function.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.