



### Synonym

IGF1R,IGFR,JTK13,CD221,MGC142170,MGC142172,MGC18216

### Source

Rat IGF-I R, His Tag(IGR-R5224) is expressed from human 293 cells (HEK293). It contains AA Glu 31 - His 936 (Accession # [P24062-1](#)).  
 Predicted N-terminus: Glu 31

### Molecular Characterization

IGF-I R(Glu 31 - His 936)  
 P24062-1 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

This protein contains a furin convertase cleavage site, 738-RRRR-741, and will be partially processed into N ( $\alpha$  chain) and C-terminal fragment (partial  $\beta$  chain) with calculated MW of 81.3 kDa and 24.2 kDa respectively. The protein migrates as 45-50 kDa (partial  $\beta$  chain), 96-115 kDa ( $\alpha$  chain) and 120 kDa ( $\alpha$  chain & partial  $\beta$  chain) under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per  $\mu$ g by the LAL method.

### Purity

>90% as determined by SDS-PAGE.

### Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at  $-20^{\circ}\text{C}$  or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- $-20^{\circ}\text{C}$  to  $-70^{\circ}\text{C}$  for 12 months in lyophilized state;
- $-70^{\circ}\text{C}$  for 3 months under sterile conditions after reconstitution.


### SDS-PAGE



Rat IGF-I R, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

### Bioactivity-ELISA

### SEC-MALS




The purity of Rat IGF-I R, His Tag (Cat. No. IGR-R5224) is more than 85% and the molecular weight of this protein is around 254-310 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,  
and more!





Immobilized Rat IGF-I R, His Tag (Cat. No. IGR-R5224) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Human IGF-I, Fc Tag with a linear range of 5-39 ng/mL (QC tested).

## Background

The Insulin-like Growth Factor 1 Receptor (IGF1) is also known as CD221, JTK13, and is a transmembrane receptor that is activated by IGF-1 and by the related growth factor IGF-2. It belongs to the large class of tyrosine kinase receptors. This receptor mediates the effects of IGF-1, which is a polypeptide protein hormone similar in molecular structure to insulin. IGF1R is made up of two alpha subunits and two beta subunits, the Both the  $\alpha$  and  $\beta$  subunits are synthesized from a single mRNA precursor. The precursor is then glycosylated, proteolytically cleaved, and crosslinked by cysteine bonds to form a functional transmembrane  $\alpha\beta$  chain. The  $\alpha$  chains are located extracellularly while the  $\beta$  subunit spans the membrane and are responsible for intracellular signal transduction upon ligand stimulation. IGF1R have a binding site for ATP, which is used to provide the phosphates for autophosphorylation. There is a 60% homology between IGF1R and the insulin receptor. In response to ligand binding, the  $\alpha$  chains induce the tyrosine autophosphorylation of the  $\beta$  chains. This event triggers a cascade of intracellular signaling that, while somewhat cell type specific, often promotes cell survival and cell proliferation.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.

Discounts, Gifts,  
and more!

