

Human IL-39 Protein, Fc Tag (MALS verified)

Catalog # IL9-H5259



Synonym

EBI3,IL39,IL-39EBI3,IL39p19,Interleukin-39,IL-39

Source

Human IL-39, Fc Tag(IL9-H5259) is expressed from human 293 cells (HEK293). It contains AA Arg 21 - Lys 229 & Arg 20 - Pro 189 (Accession # [Q14213-1](#) (IL-27 beta) & [Q9NPF7-1](#) (IL-23 alpha)).
Predicted N-terminus: Arg 21

Molecular Characterization

This protein carries a human IgG1 Fc tag at the C-terminus.
The protein has a calculated MW of 69.4 kDa. The protein migrates as 75-85 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.

- This product is stable after storage at:
- 20°C to -70°C for 12 months in lyophilized state;
 - 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human IL-39, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).


Bioactivity-ELISA

SEC-MALS



The purity of Human IL-39, Fc Tag (Cat. No. IL9-H5259) is more than 85% and the molecular weight of this protein is around 145-175 kDa verified by SEC-MALS.
[Report](#)





Immobilized Human IL-39, Fc Tag (Cat. No. IL9-H5259) at 1 µg/mL (100 µL/well) can bind Anti-IL23A MAb (Cat. No. IL9-H5259) with a linear range of 10-156 ng/mL (QC tested).

Background

Interleukin 39 (IL-39) is a new member of the IL-12 family and is composed of IL-23p19 and Ebi3 subunits. The two subunits, IL-23p19 and Ebi3, are shared with IL-23 and IL-27/IL-35. IL-23p19 is a four-helix bundle and has homology with IL-6 and granulocyte colony stimulating factor. Ebi3 was first identified in B lymphocytes infected with EB virus. Some studies have reported that the combination of IL-23R and gp130 forms the IL-39 receptor. At present, a research on IL-39 is still in its initial stage, and many issues have not been solved.

Clinical and Translational Updates

Discounts, Gifts,
and more!

