

SynonymIL1B,IL-1BETA,IL1F2,IL-1 β **Source**

Human IL-1 beta, Fc Tag (ILA-H525c) is expressed from human 293 cells (HEK293). It contains AA Ala 117 - Ser 269 (Accession # [P01584-1](#)).

Predicted N-terminus: Pro

Molecular Characterization

| | |
|---------------------------------|--|
| Fc(Pro 100 - Lys 330) P01857 | IL-1 beta(Ala 117 - Ser 269) P01584-1 |
|---------------------------------|--|

This protein carries a human IgG1 Fc tag at the N-terminus.

The protein has a calculated MW of 43.8 kDa. The protein migrates as 50-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 . Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE

Human IL-1 beta, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA**SEC-MALS**

The purity of Human IL-1 beta, Fc Tag (Cat. No. ILA-H525c) was more than 90% and the molecular weight of this protein is around 90-100 kDa verified by SEC-MALS.

[Report](#)



Immobilized Human IL-1 RII, His Tag (Cat. No. [IL2-H4226](#)) at 5 µg/mL (100 µL/well) can bind Human IL-1 beta, Fc Tag (Cat. No. ILA-H525c) with a linear range of 4.9-39 ng/mL (QC tested).

Background

Interleukin-1 beta (IL-1 β) is also known as catabolin, is a cytokine protein that in humans is encoded by the IL1B gene. IL-1 β precursor is cleaved by caspase 1 (interleukin 1 beta convertase). Cytosolic thiol protease cleaves the product to form mature IL-1 beta. IL1 β are structurally related polypeptides that share approximately 21% amino acid (aa) identity in human. Both proteins are produced by a wide variety of cells in response to inflammatory agents, infections, or microbial endotoxins. While IL1 α and IL1 β are regulated independently, they bind to the same receptor and exert identical biological effects. IL-1 β is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

References

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.