Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His,Avitag™&Tag Free (MALS verified)

Catalog # ILA-H82W3






Source

Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His, Avitag&Tag Free(ILA-H82W3) is expressed from human 293 cells (HEK293). It contains AA Glu 21 - Asp 239 & Leu 23 - Asn 254 (Accession # P16871-1 & P31785-1). Predicted N-terminus: Glu 21 & Leu 23

Molecular Characterization



Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein has a calculated MW of 32.2 kDa (IL-7 R alpha) & 31.2 kDa (IL-2 R gamma). The protein migrates as 43-65 kDa when calibrated against Star Ribbon Pre-stained Protein Marker under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using AvitagTM technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per μg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from $0.22~\mu m$ filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His, Avitag&Tag Free on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With Star Ribbon Pre-stained Protein Marker).

Bioactivity-ELISA

SEC-MALS



The purity of Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His,Avitag&Tag Free (Cat. No. ILA-H82W3) is more than 90% and the molecular weight of this protein is around 95-115 kDa verified by SEC-MALS. Report


Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His,Avitag™&Tag Free (MALS verified)







Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His,Avitag&Tag Free ELISA 0.2 μ g of Human IL-7, Fc Tag per well



Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His, Avitag& Tag Free Conc. (ng/mL)

Immobilized Human IL-7, Fc Tag (Cat. No. IL7-H5253) at 2 μ g/mL (100 μ L/well) can bind Biotinylated Human IL-7 R alpha & IL-2 R gamma Protein, His,Avitag&Tag Free (Cat. No. ILA-H82W3) with a linear range of 0.1-2 ng/mL (QC tested).

Background

The protein encoded by this gene is a receptor for interleukin 7 (IL7). The function of this receptor requires the interleukin 2 receptor, gamma chain (IL2RG), which is a common gamma chain shared by the receptors of various cytokines, including interleukins 2, 4, 7, 9, and 15. This protein has been shown to play a critical role in V(D)J recombination during lymphocyte development. Defects in this gene may be associated with severe combined immunodeficiency (SCID). Alternatively spliced transcript variants have been found.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.

