



### Synonym

IL4R,CD124,IL4RA

### Source

Cynomolgus IL-4 R alpha, His Tag (ILR-C52H8) is expressed from human 293 cells (HEK293). It contains AA Met 26 - Arg 232 (Accession # [G7Q0S7](#)).

Predicted N-terminus: Met 26

### Molecular Characterization

IL-4 R alpha(Met 26 - Arg 232)  
G7Q0S7 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 25.6 kDa. The protein migrates as 40-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per µg by the LAL method.

### Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

### Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


### SDS-PAGE



Cynomolgus IL-4 R alpha Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

### Bioactivity-ELISA

### SEC-MALS



The purity of Cynomolgus IL-4 R alpha Protein, His Tag (Cat. No. ILR-C52H8) is more than 90% and the molecular weight of this protein is around 36-49 kDa verified by SEC-MALS.

[Report](#)


Discounts, Gifts,  
and more!





## Cynomolgus IL-4 R alpha / CD124 Protein, His Tag (MALS verified) ELISA

0.5 µg of Human IL-4, premium grade per well



Immobilized Human IL-4, premium grade (Cat. No. IL4-H4218) at 5 µg/mL (100 µL/well) can bind Cynomolgus IL-4 R alpha, His Tag (Cat. No. ILR-C52H8) with a linear range of 39-312 ng/mL (QC tested).

## Background

IL-4 is a pleiotropic cytokine produced by activated Th2 cells and mast cells, and plays a pivotal role in immune responses. The effects of IL-4 are mediated after binding to high affinity receptor complexes present on hematopoietic as well as non-hematopoietic cells. Hematopoietic cellular responses to IL-4 are mediated by a high affinity receptor complex comprised of the 140 kDa IL4R $\alpha$  (CD124) subunit and the 70 kDa common cytokine  $\gamma$ c chain (CD132).

Interleukin 4 Receptor (IL4R) also known as CD124, IL4R $\alpha$  and BSF receptor, is a type I cytokine receptor produced by activated Th2 cells and mast cells, and plays an important role in Th2-biased immune responses, alternative macrophage activation, mucosal immunity, allergic inflammation, tumor progression, and atherogenesis. A soluble form of the encoded IL4R protein can be produced by an alternate splice variant or by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. IL4R can alternatively associate with IL-13Ra1 to form the type II receptor which is responsive to both IL4 and IL13. Interleukin-4 receptor has been shown to interact with SHC1.

## Clinical and Translational Updates

Discounts, Gifts,  
and more!

