



Synonym

IMD67, IPD1, IRAK-4, NY-REN-64, REN64

Source

Human IRAK4 Protein, His Tag(IR4-H5546) is expressed from Baculovirus-Insect cells. It contains AA Met 1 - Ser 460 (Accession # [Q9NWZ3-1](#)).

Predicted N-terminus: Met 1

Molecular Characterization

IRAK4(Met 1 - Ser 460)
Q9NWZ3-1 Poly-his

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 53.4 kDa. The protein migrates as 60-65 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Supplied as 0.2 µm filtered solution in 50 mM Tris, 150 mM NaCl, 1 mM DTT, pH7.5 with glycerol as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping cost.


Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.

SDS-PAGE



Human IRAK4 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Background

This gene encodes a kinase that activates NF-kappaB in both the Toll-like receptor (TLR) and T-cell receptor (TCR) signaling pathways. The protein is essential for most innate immune responses. Mutations in this gene result in IRAK4 deficiency and recurrent invasive pneumococcal disease. Multiple transcript variants encoding different isoforms have been found for this gene.

Clinical and Translational Updates

Discounts, Gifts,
and more!

