Catalog # MM8-M52H7



### Synonym

CLG1 Protein, HNC Protein, MMP-8 Protein, PMNL-CL Protein,

### Source

Mouse MMP-8 Protein, His Tag(MM8-M52H7) is expressed from human 293 cells (HEK293). It contains AA Phe 21 - Gly 262 (Accession # <u>070138-1</u>). Predicted N-terminus: Phe 21

# **Molecular Characterization**

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 28.9 kDa. The protein migrates as 35-40 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per  $\mu$ g by the LAL method.

## Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

### Formulation

Supplied as 0.2  $\mu$ m filtered solution in 25 mM MES, 150 mM NaCl, 20% Glycerol, pH 5.5 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Shipping

*This product is supplied and shipped with dry ice, please inquire the shipping cost.* 


#### Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.

### **SDS-PAGE**




Mouse MMP-8 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

# Bioactivity

Measured by its ability to cleave a fluorogenic peptide substrate Mca-KPLGL-

# SEC-MALS



The purity of Mouse MMP-8 Protein, His Tag (Cat. No. MM8-M52H7) is more than 90% and the molecular weight of this protein is around 28-40 kDa verified by SEC-MALS. <u>Report</u>

Dpa-AR-NH2. The specific activity is >700 pmol/min/µg (QC tested).

# Background



>>> www.acrobiosystems.com

4/3/2024



Catalog # MM8-M52H7

Matrix metalloproteinase-8 (MMP-8) also known as neutrophil collagenase and CLG1, is a member of matrix metalloproteinases (MMPs) family, which degrade components of the extracellular matrix (ECM) and play essential roles in various physiological processes as well as pathological processes. MMP-8 may affect the metastatic behavior of breast cancer cells through protection against lymph node metastasis, underlining the importance of anti-target identification in drug development. MMP-8 in the tumor may have a protective effect against lymph node metastasis.

### **Clinical and Translational Updates**



>>> www.acrobiosystems.com

