

### Synonym

NCR1,LY94,CD335,NK-p46,hNKp46

#### Source

Human NKp46, His Tag(NC1-H52H6) is expressed from human 293 cells (HEK293). It contains AA Gln 22 - Arg 258 (Accession # <u>076036-1</u>).

## **Molecular Characterization**

NKp46(Gln 22 - Arg 258) O76036-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 28.9 kDa. The protein migrates as 35-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### **Endotoxin**

Less than 1.0 EU per µg by the LAL method.

## **Purity**

>90% as determined by SDS-PAGE.

#### **Formulation**

Lyophilized from 0.22  $\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

#### **Storage**


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

### **SDS-PAGE**




Human NKp46, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

## **Bioactivity-ELISA**

# Human NKp46 / NCR1 / CD335 Protein, His Tag

Catalog # NC1-H52H6





Immobilized Human NKp46, His Tag (Cat. No. NC1-H52H6) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Anti-NKP46 Antibody, Human IgG1 with a linear range of 0.1-2 ng/mL (QC tested).

# **Background**

Natural cytotoxicity triggering receptor 1 (NCR1) is also known as Natural killer cell p46-related protein (NK-p46), Lymphocyte antigen 94 homolog (LY94), CD antigen CD335, which belongs to the natural cytotoxicity receptor (NCR) family. NCR1 contains two Ig-like (immunoglobulin-like) domains. NCR1 interacts with CD247 and FCER1G. NCR1 / CD335 may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis.

# **Clinical and Translational Updates**

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.