

Synonym

PDCD1,PD1,CD279,SLEB2

Source

Rabbit PD-1, Fc Tag (PD1-RB5251) is expressed from human 293 cells (HEK293). It contains AA Leu 25 - Gln 167 (Accession # G1SUF0).

Predicted N-terminus: Leu 25

Molecular Characterization

PD-1(Leu 25 - Gln 167) G1SUF0	Fc(Pro 100 - Lys 330) P01857
----------------------------------	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 42.6 kDa. The protein migrates as 55-66 kDa under reducing (R) condition (SDS-PAGE).

Endotoxin

Less than 1.0 EU per μg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Rabbit PD-1, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Immobilized Mouse PD-L1, His Tag (Cat. No. PD1-M5220) at 10 µg/mL (100 µL/well) can bind Rabbit PD-1, Fc Tag (Cat. No. PD1-RB5251) with a linear range of 3-50 ng/mL (QC tested).

Background

Programmed cell death protein 1 (PD-1) is also known as CD279 and PDCD1, is a type I membrane protein and is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 is expressed on the surface of activated T cells, B cells, macrophages, myeloid cells and a subset of thymocytes. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. PD-L1 is expressed on almost all murine tumor cell lines, including PA1 myeloma, P815 mastocytoma, and B16 melanoma upon treatment with IFN-γ. PD-L2 expression is more restricted and is expressed mainly by DCs and a few tumor lines. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN-γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediated signal by dephosphorylating key signal transducer. In vitro, treatment of anti-CD3 stimulated T cells with PD-L1-Ig results in reduced T cell proliferation and IFN-γ secretion. Monoclonal antibodies targeting PD-1 that boost the immune system are being developed for the treatment of cancer.

References

- (1) [Ishida Y., et al., 1992, EMBO J. 11 \(11\): 3887-95.](#)
- (2) [Blank C., et al., 2007, Cancer Immunol. Immunother. 56 \(5\): 739-45.](#)
- (3) [Agata Y., et al., 1996, Int. Immunol. 8 \(5\): 765-72.](#)
- (4) [Freeman GJ., et al., 2000, J. Exp. Med. 192 \(7\): 1027-34.](#)
- (5) [Latchman Y., et al., 2001, Nat. Immunol. 2 \(3\): 261-8.](#)
- (6) [Yamazaki T., et al., 2002, J. Immunol. 169 \(10\): 5538-45.](#)

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.