

Synonym

C7orf15,C7orf15MGC138295,CD112R,MGC104322,MGC138297,MGC2463,P VRIG,CD112 receptor

Source

Human PVRIG, Fc Tag (PVG-H5257) is expressed from human 293 cells (HEK293). It contains AA Thr 41 - Asp 171 (Accession # Q6DK17-1).

Molecular Characterization

PVRIG(Thr 41 - Asp 171) Fc(Pro 100 - Lys 330)
Q6DKI7-1 P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 40.3 kDa. The protein migrates as 45-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from $0.22 \mu m$ filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20° C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human PVRIG, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS




The purity of Human PVRIG, Fc Tag (Cat. No. PVG-H5257) is more than 90% and the molecular weight of this protein is around 77-105 kDa verified by SEC-MALS.

Report

Human PVRIG Protein, Fc Tag (MALS verified)







Immobilized Human Nectin-2, His Tag (Cat. No. <u>PV2-H52E2</u>) at 5 μ g/mL (100 μ L/well) can bind Human PVRIG, Fc Tag (Cat. No. <u>PVG-H5257</u>) with a linear range of 5-78 ng/mL (QC tested).

Background

Human PVRIG (poliovirus receptor related immunoglobulin domain-containing protein), also known as CD112 receptor (CD112R), is an approximately 34 kDa single transmembrane protein in the poliovirus receptor-like protein (PVR) family. The CD112R gene encodes a putative single transmembrane protein, which is composed of a single extracellular IgV domain, one transmembrane domain, and a long intracellular domain. Notably, the intracellular domain of phatases. The extracellular domain sequence of human and mouse CD112R have 65.3% similarity. CD112R may act as a coinhibitory receptor that suppresses T-cell receptor-mediated signals.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.