


Synonym

S1 protein NTD,Spike protein S1 NTD,BetaCoV S1-NTD

Source

Biotinylated SARS-CoV-2 S1 protein NTD, His,Avitag (S1D-C52E2) is expressed from human 293 cells (HEK293). It contains AA Val 16 - Phe 318 (Accession # [QHD43416.1](#)).
Predicted N-terminus: Val 16

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 38.0 kDa. The protein migrates as 53-90 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Biotinylation

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Biotin:Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.
>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:
• -20°C to -70°C for 12 months in lyophilized state;
• -70 °C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated SARS-CoV-2 S1 protein NTD, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.


Bioactivity-ELISA

SEC-MALS



The purity of Biotinylated SARS-CoV-2 S1 protein NTD, His,Avitag (Cat. No. S1D-C52E2) is more than 90% and the molecular weight of this protein is around 70-80 kDa verified by SEC-MALS.

[Report](#)



Immobilized Anti-SARS-CoV-2 Spike NTD Antibody, Chimeric mAb (Cat. No. [SPD-M121](#)) at 1 µg/mL (100 µL/well) can bind Biotinylated SARS-CoV-2 S1 protein NTD, His,Avitag (Cat. No. [S1D-C52E](#)) with a linear range of 1-25 ng/mL (QC tested).

Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.