

Synonym


Spike,S1 protein,Spike glycoprotein Subunit1,Spike protein S1

Source

Biotinylated SARS-CoV-2 S1 protein (D614G), His,Avitag(S1N-C82E3) is expressed from human 293 cells (HEK293). It contains AA Val 16 - Arg 685 (Accession # QHD43416.1 (D614G)).

Predicted N-terminus: Val 16

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (AvitagTM)

The protein has a calculated MW of 78.5 kDa. The protein migrates as 100-115 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using AvitagTM technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Biotinylated SARS-CoV-2 S1 protein (D614G), His, Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

Biotinylated SARS-CoV-2 (COVID-19) S1 protein (D614G), His,Avitag™


Catalog # S1N-C82E3





Immobilized Human ACE2, Fc Tag (Cat. No. AC2-H5257) at 1 μ g/mL (100 μ L/well) can bind Biotinylated SARS-CoV-2 S1 protein (D614G), His,Avitag (Cat. No. S1N-C82E3) with a linear range of 0.02-0.625 μ g/mL (QC tested).

Biotinylated SARS-CoV-2 S1 protein (D614G), His,Avitag ELISA 0.1 μ g of Biotinylated SARS-CoV-2 S1 protein (D614G), His,Avitag per well



Anti-SARS-CoV-2 Neutralizing Antibody, Human IgG1 Conc. (ng/mL)

Immobilized Biotinylated SARS-CoV-2 S1 protein (D614G), His,Avitag (Cat. No. S1N-C82E3) at 1 μ g/mL (100 μ L/well) can bind Anti-SARS-CoV-2 Neutralizing Antibody, Human IgG1 (Cat. No. SAD-S35) with a linear range of 0.8-6 ng/mL (Routinely tested).

Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.