

Source

Anti-SARS-CoV-2 Spike RBD Neutralizing Antibody, Mouse IgG1 (AM122) (SCT-M369), originally from mouse immunized with recombinant SARS-CoV-2 Spike RBD protein, is recombinantly expressed from HEK293 cells.

Isotype

Mouse IgG1/kappa

Specificity

This product is a specific antibody against SARS-CoV-2 Spike protein RBD domain. Cross-reactivity with Spike protein RBD domain of other coronaviruses, including SARS-CoV, MERS-CoV, HCoV-229E, HCoV-NL63, HCoV-OC43 and HCoV-HKU1 has not been tested.

Purity

>95% as determined by SDS-PAGE.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Formulation

Lyophilized from $0.22~\mu m$ filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Anti-SARS-CoV-2 Spike RBD Neutralizing Antibody, Mouse IgG1 (AM122) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-Elisa

Anti-SARS-CoV-2 Spike RBD Neutralizing Antibody, Mouse IgG1 (AM122)







Immobilized SARS-CoV-2 Spike Trimer, His Tag (B.1.1.529/Omicron) (Cat. No. SPN-C52Hz) at $2\mu g/mL$ ($100\mu L/well$) can bind Anti-SARS-CoV-2 Spike RBD Neutralizing Antibody, Mouse IgG1 (AM122) (Cat. No. SCT-M369) with a linear range of 0.2-6 ng/mL (QC tested).

Background

Its been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Clinical and Translational Updates

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.