

### Synonym

Spike, Sprotein, Spike glycoprotein, Sglycoprotein

#### Source

Biotinylated SARS-CoV-2 S protein (D614G), His,Avitag, Super stable trimer (SPN-C82E3) is the ectodomain of SARS-CoV-2 S protein that contains AA Val 16 - Pro 1213 (Accession # QHD43416.1 (D614G)) and D614G mutation, which has become increasingly common in SARS-CoV-2 viruses from around the world . The recombinant protein is expressed from human 293 cells (HEK293) with T4 fibritin trimerization motif and a polyhistidine tag at the C-terminus. Proline substitutions (F817P, A892P, A899P, A942P, K986P, V987P) and alanine substitutions (R683A and R685A) are introduced to stabilize the trimeric prefusion state of SARS-CoV-2 S protein and abolish the furin cleavage site, respectively.

Predicted N-terminus: Val 16

#### **Molecular Characterization**

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag<sup>TM</sup>)

The protein has a calculated MW of 139.7 kDa. The protein migrates as 150-200 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### Labeling

Biotinylation of this product is performed using Avitag<sup>TM</sup> technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

### **Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

# Endotoxin

Less than  $1.0 \ EU$  per  $\mu g$  by the LAL method.

### **Purity**

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

#### **Formulation**

Lyophilized from 0.22 µm filtered solution in PBS with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

### Storage


For long term storage, the product should be stored at lyophilized state at -20 $^{\circ}$ C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## **SDS-PAGE**



Biotinylated SARS-CoV-2 S protein (D614G), His, Avitag, Super stable trimer on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

## **SEC-MALS**




The purity of Biotinylated SARS-CoV-2 S protein (D614G), His,Avitag, Super stable trimer (Cat. No. SPN-C82E3) is more than 90% and the molecular weight of this protein is around 520-620 kDa verified by SEC-MALS.

Report



### **Bioactivity-ELISA**


Biotinylated SARS-CoV-2 S protein (D614G), His, Avitag, Super stable trimer ELISA 0.1  $\mu$ g of Human ACE2, Fc Tag per well



SARS-CoV-2 S protein (D614G), His, Avitag, Super stable trimer Conc. (ng/mL)

Immobilized Human ACE2, Fc Tag (Cat. No. AC2-H5257) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Biotinylated SARS-CoV-2 S protein (D614G), His,Avitag, Super stable trimer (Cat. No. SPN-C82E3) with a linear range of 2-63 ng/mL (QC tested).

SARS-CoV-2 S protein (D614G), His, Avitag, Super stable trimer ELISA 0.1 µg of SARS-CoV-2 S protein (D614G), His, Avitag, Super stable trimer per well



Anti-SARS-CoV-2 RBD Neutralizing Antibody, Human IgG1 Conc. (ng/mL)

Immobilized Biotinylated SARS-CoV-2 S protein (D614G), His,Avitag, Super stable trimer (Cat. No. SPN-C82E3) at 1 μg/mL (100 μL/well) on Recombinant Streptavidin (Cat. No. STN-N5116) precoated (0.5 μg/well) plate, can bind Anti-SARS-CoV-2 RBD Neutralizing Antibody, Human IgG1 (Cat. No. SAD-S35) with a linear range of 0.1-2 ng/mL (Routinely tested).

## Background

It's been reported that SARS-CoV-2 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

# **Clinical and Translational Updates**

Please contact us via <u>TechSupport@acrobiosystems.com</u> if you have any question on this product.