



Synonym

Streptavidin,SA

Source

Streptavidin Protein-Alexa Fluor 647(STN-NA119) is expressed from E. coli cells.

Molecular Characterization

This protein carries no "tag".

The protein has a calculated MW of 13.8 kDa. The protein migrates as 12-15 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE).

Conjugate

AF647

Excitation Wavelength: 640 nm

Emission Wavelength: 672 nm

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with AF647 using standard chemical labeling method. The residual AF647 is removed by molecular sieve treatment during purification process.

Protein Ratio

The AF647 to protein molar ratio is 3-6.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-HPLC.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Streptavidin Protein-Alexa Fluor 647 on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-HPLC




The purity of Streptavidin Protein-Alexa Fluor 647 (Cat. No. STN-NA119) was greater than 90% as determined by SEC-HPLC.

Bioactivity-FACS

Discounts, Gifts,
and more!





5e5 of Jurkat cells were stained with 100 μ L of 3 μ g/mL of Biotinylated Anti-Human CD3 Antibody, Mouse IgG2a, Avitag (Clone: OKT3), premium grade (Cat. No. CDE-M525a) and isotype control antibody respectively, washed and then followed with 100 μ L of 1 μ g/mL of Streptavidin Protein-Alexa Fluor 647 (Cat. No. STN-NA119) and analyzed with FACS. Alexa 647 signal was used to evaluate the binding activity (Routinely tested).

Background

Streptavidin is a tetrameric protein purified from the bacterium *Streptomyces avidinii*, and exhibits high binding affinity for biotin. Able to bind one molecule of biotin with each subunit. Streptavidin (PI=6.0-7.5) has lower level of non-specific binding to various biological components at physiological pH than avidin (PI=7.4), resulting from its isoelectric point (PI). Streptavidin is useful in affinity chromatography, ELISA, immunohistochemistry and Western Blotting.

Clinical and Translational Updates

Discounts, Gifts,
and more!

