


Synonym

CD71,TFRC,TFR,,TfR1,Trfr,T9,p90

Source

Biotinylated Cynomolgus Transferrin R, His Tag, primary amine labeling (TFR-C8249) is expressed from human 293 cells (HEK293). It contains AA Cys 89 - Phe 760 (Accession # [G8F602-1](#)). It is the biotinylated form of Cynomolgus Transferrin R, His Tag(Cat. No. TFR-C524a).
Predicted N-terminus: His

Molecular Characterization



This protein carries a polyhistidine tag at the N-terminus
The protein has a calculated MW of 77.1 kDa. The protein migrates as 70-90 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with biotins using standard chemical labeling method. A standard biotin reagent (13.5 angstroms) is used in this product.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Biotinylated Cynomolgus Transferrin R, His Tag, primary amine labeling on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Immobilized Biotinylated Cynomolgus Transferrin R, His Tag, primary amine labeling (Cat. No. TFR-C8249) at 1 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate can bind Mouse Transferrin, Mouse IgG2a Fc Tag, low endotoxin (Cat. No. TRN-M525b) with a linear range of 0.3-10 ng/mL (QC tested).



Immobilized Mouse Transferrin, Mouse IgG2a Fc Tag, low endotoxin (Cat. No. TRN-M525b) at 2 µg/mL (100 µL/well) can bind Biotinylated Cynomolgus Transferrin R, His Tag, primary amine labeling (Cat. No. TFR-C8249) with a linear range of 0.3-10 ng/mL (Routinely tested).

Background

CD antigen CD71 is also known as Transferrin receptor protein 1, TfR, sTfR, p90, TfR1, Trfr, which belongs to the peptidase M28 family and M28B subfamily. CD71 /TfR contains one PA (protease associated) domain. CD71 / TfR1 is required for iron delivery from transferrin to cells. CD71 is a potential new target in cases of human leukomia & lymphoma. CD71 /TFRC / TfR has been shown to interact with GABARAP and HFE.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.