


SynonymTGFBR2,TGFR2,TbetaR-II,TGF β R2**Source**

Biotinylated Human TGF-beta RII, His,Avitag (TG2-H82E4) is expressed from human 293 cells (HEK293). It contains AA Thr 23 - Asp 159 (Accession # [P37173-1](#)).

Predicted N-terminus: Thr 23

Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 19.1 kDa. The protein migrates as 28-40 kDa under reducing (R) condition (SDS-PAGE) due to Glycosylation.

Biotinylation

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Biotin:Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per μ g by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.



This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Biotinylated Human TGF-beta RII, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA



Immobilized Human Latent TGFB1, His Tag (Cat. No. [TG1-H524x](#)) at 1 µg/mL (100 µL/well) can bind Biotinylated Human TGF-beta RII, His,Avitag (Cat. No. [TG2-H82E4](#)) with a linear range of 5-78 ng/mL (QC tested).

Serial dilutions of Human TGF-beta RII, His Tag (Cat. No. [TG2-H52H5](#)) were added into Human Latent TGFB1, His Tag (Cat. No. [TG1-H524x](#)) : Biotinylated Human TGF-beta RII, His,Avitag (Cat. No. TG2-H82E4) binding reactions. The half maximal inhibitory concentration (IC₅₀) is 0.39287 µg/mL (Routinely tested).

Background

TGF-beta receptor type-2 (TGFR2 or TGFR-2) is also known as TGF-beta type II receptor, Transforming growth factor-beta receptor type II, TbetaR-II, TGF β R2, which is a homodimer or heterohexamer, belongs to the protein kinase superfamily, TKL Ser/Thr protein kinase family and TGFB receptor subfamily. TGFR2 / TGFR2 binds TGF- β 1 / TGFB1 and TGF- β 3 / TGFB3 with high affinity and TGF- β 2 / TGFB2 with a much lower affinity. This type II receptor forms a heterodimeric complex with type I receptor and is essential for signal transduction. Upon ligand binding, the TGFR2 autophosphorylates its cytoplasmic domain and subsequently phosphorylates the downstream molecules which then enter the nucleus and regulate the transcription of a subset of genes related to cell proliferation.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.