Catalog # TR2-M82F3



Synonym

TACSTD2,GA733-1,M1S1,TROP2


Source

Biotinylated Mouse TROP-2 Protein, Fc, Avitag(TR2-M82F3) is expressed from human 293 cells (HEK293). It contains AA Gln 25 - Gly 270 (Accession #

<u>Q8BGV3-1</u>).

Predicted N-terminus: Gln 25

Molecular Characterization



This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (AvitagTM).

The protein has a calculated MW of 56.1 kDa. The protein migrates as 75-90 kDa when calibrated against Star Ribbon Pre-stained Protein Marker under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag[™] technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.


Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per μ g by the LAL method.

SDS-PAGE



Biotinylated Mouse TROP-2 Protein, Fc, Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.



The purity of Biotinylated Mouse TROP-2 Protein, Fc, Avitag (Cat. No. TR2-M82F3) is more than 90% and the molecular weight of this protein is around 120-145 kDa verified by SEC-MALS.

protein is greater than 90% (With Star Ribbon Pre-stained Protein Marker).

Report

Background

>> www.acrobiosystems.com

9/7/2023



Catalog # TR2-M82F3

TROP-2 is a single-copy gene in human cells, and encodes a type-1 transmembrane glycoprotein which is over-expressed in various malignancies, also referred to as tumor associated calcium signal transducer 2 (TACSTD2), GA733-1 or M1S1. TROP-2 is related to epithelial cell adhesion molecule (EpCAM), also called TROP-1, gp40, and KSA. Trop-1 and Trop-2 are homologous to serum IGF-II-binding proteins and appear as signal transducers. Thus, they likely represent novel cell-surface receptors and may play a role in regulating the growth of carcinoma cells.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.



9/7/2023